

Farmcell Usage Schedule for CORN



	Before Plowing	After Plowing	Seed Soaking (15 to 30min)	One week after sowing	Day 30	Day 60	Ripening stage
Dilution	1 : 150	1 : 150	1 : 300	1 : 500	1 : 400	1 : 300	
Farmcell Product							
Usage of Farmcell with 20L water	135ml	135ml	67ml	40ml	50ml	67ml	

Mixing Ratio Reference: Farmcell / Water
 Pump: 135ml : 20L Water
 Drone : 400ml : 60L Water

Reference Farmcell 1.0L / Hectare or
 usage per Land size: 0.4L / Acre

Precaution:

1. Use water without chlorine (Non-Chlorine Water).
2. Do not mix with pesticides or fertilizers.
3. Spray equipment must be used separately.

山东康耕农业发展有限公司

玉米- 植物干细胞使用方法



	翻土前	翻土后	泡种子 (15 至 30分钟)	播种一周后	第 30天	第 60天	待收割
稀释比率	1 : 150	1 : 150	1 : 300	1 : 500	1 : 400	1 : 300	
植物干细胞 产品							
稀释比率 用量 (兑 20公升水)	135毫升	135毫升	67毫升	40毫升	50毫升	67毫升	

喷施方式: 叶面喷洒/根灌

使用范围参考: 1公升FARMCELL 对 1公顷地 (15亩)

1. 使用无氯水 (Non-Chlorine Water) 。
2. 请勿与其他农药或肥料复配使用。
3. 喷射器材必须独立使用, 喷洒过农药的器材不可混用。
4. 稀释后的原液必须在两天内使用。
5. 养地时添加有机底肥, 例如有机鸡粪肥或固有的方式。