



TRIAL SOP

FARMCELL® PESTACURE

For Rubber Pestalotiopsis Disease

MATERIALS:

1. FARMCELL GDS+
2. FARMCELL PestaCure



Phase 1: Spray continuously once a week for 3 weeks

Duration	MATERIAL	APPLICATION	DILUTION
0 ~ 3 weeks	FARMCELL GDS+	Spray surrounding the tree with a diameter of approximately 2.0 meters	1:50
	FARMCELL PestaCure	Spray on trunk approximately 2.0 meters height	1:50

Phase 2: Spray consistently every two weeks for a duration of two months.

Duration	MATERIAL	APPLICATION	DILUTION
Start from 5 th week	FARMCELL GDS+	Spray around the roots with a diameter of approximately 2.0 meters	1:50
	FARMCELL PestaCure	Spray on trunk approximately 2.0 meters height	1:50

Usage steps: Apply GDS+ first, then apply PestaCure after 30 minutes.

Remark: FARMCELL GDS+ Only spray on Ground.

Phase 3: Spray consistently every 2 months

Duration	MATERIAL	APPLICATION	DILUTION
Every 2 Months	FARMCELL GDS+	Spray around the roots with a diameter of approximately 2.0 meters	1:50
	FARMCELL PestaCure	Spray on trunk approximately 2.0 meters height	1:150

Precaution:

1. Use non-chlorine water ONLY.
2. Do not mix with other pesticides or fertilizers.
3. The diluted solution must be used within two days
4. Spraying equipment must be used independently, and equipment that has been used to spray pesticides cannot be mixed.